

Application of the generalized Kudryashov method to the Eckhaus equation

Ahmed H. Arnous^a, Mohammad Mirzazadeh^b

^aDepartment of Engineering Mathematics and Physics,
Higher Institute of Engineering,
El Shorouk, Egypt
ahmed.h.arnous@gmail.com

^bDepartment of Mathematics, Faculty of Mathematical Sciences,
University of Guilan,
Rasht, Iran
mirzazadehs2@guilan.ac.ir

Received: October 29, 2014 / **Revised:** June 5, 2015 / **Published online:** June 23, 2016

Abstract. In this paper, the generalized Kudryashov method is presented to seek exact solutions of the Eckhaus equation. From these solutions we can derive solitary wave solutions as a special case. The proposed method is direct, effective and convenient and can be applied to many nonlinear evolution equations in mathematical physics.

Keywords: nonlinear evolution equations, exact solutions, Eckhaus equation, generalized Kudryashov method.

1 Introduction

It is well known that nonlinear evolution equations (NLEEs) are widely used to describe physical phenomena in various scientific and engineering fields, such as fluid mechanics, plasma physics, optical fibers, biology, solid state physics, etc. [1–7, 10, 12–15, 20, 21, 23–29, 31–36]. In order to understand the mechanisms of those physical phenomena, it is necessary to explore their solutions and properties. Solutions for the NLEEs can not only describe the designated problems, but also give more insights on the physical aspects of the problems in the related fields. In recent years, various powerful methods have been presented for finding exact solutions of the NLEEs in mathematical physics, such as tanh-function method [28], extended tanh-function method [10, 29], sine-cosine method [27], Jacobi elliptic function method [13, 32], F -expansion method [1, 31], exp-function method [14], (G'/G) -expansion method [26], Q -function method [20] and so on.

The Q -function method, which is a direct and effective algebraic method for computing exact travelling wave solutions, was first proposed by Kudryashov [16]. The Q -function method that is known as the Kudryashov method is one of the most effective methods

for finding the exact solution of high order NLEEs [17]. The most complete description of this method was given in [19]. The successful application of this method to NLEEs was performed in works [8, 18, 22]. In the present work, we apply the generalized Kudryashov method [8] to the Eckhaus equation, which has the following form:

$$i\psi_t + \psi_{xx} + 2(|\psi|^2)_x \psi + |\psi|^4 \psi = 0,$$

where $\psi = \psi(x, t)$, $\psi : \mathbb{R}^2 \rightarrow \mathbb{C}$ is a complex-valued function of two real variables x, t . This equation is a nonlinear Schrödinger-type equation (NLSE) that can be linearized to the free linear Schrödinger equation, which was found in [9] as an asymptotic multiscale reduction of certain classes of nonlinear partial differential equations. In [11], many of the properties of the Eckhaus equation were investigated, including the linearization, soliton solutions etc.

This NLSE is another nonlinear evolution equation that is available in the literature. In fact, this is a dissipative equation. Therefore, this is not studied in the context of fiber optics. The first term is the linear evolution term, while the second term is accounted for dispersion that is commonly referred to as group velocity dispersion (GVD). The third term is a dissipative terms that is responsible for the damping of the soliton solution. Finally, the last term is the nonlinear term.

2 The generalized Kudryashov method

Suppose that we have a nonlinear evolution equation in the form

$$F(u, u_t, u_x, u_{xx}, u_{xt}, \dots) = 0, \quad (1)$$

where $u = u(x, t)$ is an unknown function, F is a polynomial in u and its various partial derivatives u_t, u_x with respect to t, x respectively, in which the highest order derivatives and nonlinear terms are involved.

Step 1. Using the traveling wave transformation

$$u(x, t) = u(\xi), \quad \xi = k(x - ct) + \xi_0, \quad (2)$$

where ξ_0 is an arbitrary constant and k, c are constant to be determined later. Then Eq. (1) is reduced to a nonlinear ordinary differential equation (NODE) of the form

$$P(u, u_\xi, u_{\xi\xi}, \dots) = 0. \quad (3)$$

Step 2. Suppose that the solution of Eq. (3) has the following form:

$$u(\xi) = \frac{\sum_{i=0}^N a_i Q^i(\xi)}{\sum_{j=0}^M b_j Q^j(\xi)} = \frac{A(Q(\xi))}{B(Q(\xi))}, \quad (4)$$

where a_i ($i = 0, 1, \dots, N$) and b_j ($j = 0, 1, \dots, M$) are constants to be determined such that $a_N \neq 0, b_M \neq 0$ and

$$Q(\xi) = \frac{1}{1 + \kappa e^\xi} \quad (5)$$

is the solution of the equation

$$Q_\xi = Q^2 - Q, \quad (6)$$

where κ is an arbitrary constant.

Step 3. Determine the positive integer numbers N and M in Eq. (4) by using the homogeneous balance between the highest order derivatives and the nonlinear terms in Eq. (3) after substituting Eq. (6) and the necessary derivatives of u , which have the form

$$u_\xi = (Q^2 - Q) \left(\frac{A'B - AB'}{B^2} \right), \quad (7)$$

$$u_{\xi\xi} = \frac{(Q^2 - Q)^2}{B^3} \{ B(BA'' - AB'') - 2B'(A'B - AB') \} \\ + (2Q - 1)(Q^2 - Q) \left(\frac{A'B - AB'}{B^2} \right), \quad (8)$$

where the prime ' denotes the derivative d/dQ .

Step 4. Substitute Eqs. (4), (7) and (8) into Eq. (3). As a result of this substitution, we get a polynomial of Q . In this polynomial we gather all terms of same powers and equating them to be zero, we obtain a system of algebraic equations, which can be solved by the Maple or Mathematica to get the unknown parameters a_i ($i = 0, 1, \dots, N$), b_j ($j = 0, 1, \dots, M$), k , c . Consequently, we obtain the exact solutions of Eq. (1).

3 The Eckhaus equation

The Eckhaus equation is a nonlinear Schrödinger-type equation, which can be written as

$$i\psi_t + \psi_{xx} + 2(|\psi|^2)_x \psi + |\psi|^4 \psi = 0. \quad (9)$$

This equation has been solved by using the (G'/G) -expansion method [30]. Let us now solve Eq. (9) by using the generalized Kudryashov method. To this end, we use the following wave transformation:

$$\psi(x, t) = u(\xi) e^{i(\alpha x + \beta t)}, \quad \xi = k(x - 2\alpha t) + \xi_0, \quad (10)$$

where k , α , β are constants to be determined later and ξ_0 is an arbitrary constant. Now Eq. (9) is reduced to the following NODE:

$$k^2 u_{\xi\xi} - (\beta + \alpha^2)u + 4k u_\xi u^2 + u^5 = 0. \quad (11)$$

Suppose that

$$u = v^{1/2}. \quad (12)$$

Then Eq. (11) can be reduced to the following NODE:

$$2k^2 v v_{\xi\xi} - k^2 v_\xi^2 - 4(\beta + \alpha^2)v^2 + 8k v^2 v_\xi + 4v^4 = 0. \quad (13)$$

Balancing $vv_{\xi\xi}$ with v^4 in Eq. (13), we get the formula $N = M + 1$. If we choose $M = 1$ and $N = 2$, then

$$v = \frac{A(Q)}{B(Q)} = \frac{a_0 + a_1Q + a_2Q^2}{b_0 + b_1Q}. \quad (14)$$

Substituting v and it's their necessary derivatives into (13) and equating all the coefficients of Q to zero, we obtain

$$3a_2^2b_1^2k^2 + 8a_2^3b_1k + 4a_2^4 = 0, \quad (15)$$

$$-4a_2^2b_1^2k^2 + 8a_2^2b_0b_1k^2 + 4a_1a_2b_1^2k^2 + 16a_2^3b_0k - 8a_2^3b_1k + 16a_1a_2^2b_1k + 16a_1a_2^3 = 0, \quad (16)$$

$$-4\alpha^2a_2^2b_1^2 - 4a_2^2\beta b_1^2 + 8a_2^2b_0^2k^2 + a_2^2b_1^2k^2 - 10a_2^2b_0b_1k^2 + 6a_0a_2b_1^2k^2 - 6a_1a_2b_1^2k^2 + 10a_1a_2b_0b_1k^2 - 16a_2^3b_0k + 40a_1a_2^2b_0k + 8a_0a_2^2b_1k - 16a_1a_2^2b_1k + 8a_1^2a_2b_1k + 16a_0a_2^3 + 24a_1^2a_2^2 = 0, \quad (17)$$

$$-8\alpha^2a_2a_1b_1^2 - 8\alpha^2a_2^2b_0b_1 - 8a_2a_1\beta b_1^2 - 8a_2^2\beta b_0b_1 + 12a_2a_1b_0^2k^2 + 2a_2a_1b_1^2k^2 - 12a_2a_1b_0b_1k^2 - 12a_2^2b_0^2k^2 - 12a_0a_2b_1^2k^2 + 2a_2^2b_0b_1k^2 + 12a_0a_2b_0b_1k^2 + 32a_2a_1^2b_0k - 8a_2a_1^2b_1k - 40a_2^2a_1b_0k + 32a_0a_2^2b_0k - 8a_0a_2^2b_1k + 16a_2a_1^3 + 48a_0a_2^2a_1 = 0, \quad (18)$$

$$-4\alpha^2a_1^2b_1^2 - 16\alpha^2a_2a_1b_0b_1 - 4\alpha^2a_2^2b_0^2 - 8\alpha^2a_0a_2b_1^2 - 4a_1^2\beta b_1^2 - 16a_2a_1\beta b_0b_1 - 4a_2^2\beta b_0^2 - 8a_0a_2\beta b_1^2 + 3a_1^2b_0^2k^2 + 2a_1^2b_0b_1k^2 - 18a_2a_1b_0^2k^2 - 2a_0a_1b_1^2k^2 - 2a_0a_1b_0b_1k^2 + 2a_2a_1b_0b_1k^2 + 4a_2^2b_0^2k^2 + 12a_0a_2b_0^2k^2 - a_0^2b_1^2k^2 + 6a_0a_2b_1^2k^2 - 20a_0a_2b_0b_1k^2 + 8a_1^3b_0k - 32a_2a_1^2b_0k - 8a_0a_1^2b_1k + 48a_0a_2a_1b_0k - 32a_0a_2^2b_0k - 8a_0^2a_2b_1k + 4a_1^4 + 48a_0a_2a_1^2 + 24a_0^2a_2^2 = 0, \quad (19)$$

$$-8\alpha^2a_1^2b_0b_1 - 8\alpha^2a_2a_1b_0^2 - 8\alpha^2a_0a_1b_1^2 - 16\alpha^2a_0a_2b_0b_1 - 8a_1^2\beta b_0b_1 - 8a_2a_1\beta b_0^2 - 8a_0a_1\beta b_1^2 - 16a_0a_2\beta b_0b_1 - 4a_1^2b_0^2k^2 - 2a_1^2b_0b_1k^2 + 4a_0a_1b_0^2k^2 + 6a_2a_1b_0^2k^2 + 2a_0a_1b_1^2k^2 + 4a_0a_1b_0b_1k^2 - 20a_0a_2b_0^2k^2 - 4a_0^2b_0b_1k^2 + 8a_0a_2b_0b_1k^2 - 8a_1^3b_0k + 16a_0a_1^2b_0k + 8a_0a_1^2b_1k - 48a_0a_2a_1b_0k - 16a_0^2a_1b_1k + 16a_0^2a_2b_0k + 8a_0^2a_2b_1k + 16a_0a_1^3 + 48a_0^2a_2a_1 = 0, \quad (20)$$

$$-4\alpha^2a_0^2b_1^2 - 8\alpha^2a_2a_0b_0^2 - 16\alpha^2a_1a_0b_0b_1 - 4\alpha^2a_1^2b_0^2 - 4a_0^2\beta b_1^2 - 8a_2a_0\beta b_0^2 - 16a_1a_0\beta b_0b_1 - 4a_1^2\beta b_0^2 + a_0^2b_1^2k^2 + 6a_0^2b_0b_1k^2 - 6a_1a_0b_0^2k^2 + 8a_2a_0b_0^2k^2 - 2a_1a_0b_0b_1k^2 + a_1^2b_0^2k^2 - 8a_0^3b_1k + 8a_1a_0^2b_0k - 16a_2a_0^2b_0k + 16a_1a_0^2b_1k - 16a_1^2a_0b_0k + 16a_2a_0^3 + 24a_1^2a_0^2 = 0, \quad (21)$$

$$-8\alpha^2a_0^2b_0b_1 - 8\alpha^2a_1a_0b_0^2 - 8a_0^2\beta b_0b_1 - 8a_1a_0\beta b_0^2 - 2a_0^2b_0b_1k^2 + 2a_1a_0b_0^2k^2 + 8a_0^3b_1k - 8a_1a_0^2b_0k + 16a_1a_0^3 = 0, \quad (22)$$

$$-4\alpha^2a_0^2b_0^2 - 4a_0^2\beta b_0^2 + 4a_0^4 = 0. \quad (23)$$

Solving system (15)–(23) with the aid of Mathematica, we obtain the following results.

Result 1.

$$a_0 = 0, \quad a_1 = \frac{-kb_0}{2}, \quad a_2 = \frac{-kb_1}{2}, \quad k = \pm 2\sqrt{\beta + \alpha^2}. \quad (24)$$

Substituting (24) into (14) with (5), (10) and (12), we obtain the following solutions of Eq. (9):

$$\psi_{1,2}(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{1 + \kappa e^{\pm 2\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0}} \right\}^{1/2} e^{i(\alpha x + \beta t)}, \quad (25)$$

$$\psi_{3,4}(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{1 + \kappa e^{\pm 2\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0}} \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (26)$$

If we set $\kappa = 1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 - \tanh(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}, \quad (27)$$

$$\psi(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 - \tanh(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (28)$$

If we set $\kappa = -1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 - \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}, \quad (29)$$

$$\psi(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 - \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (30)$$

Result 2.

$$a_0 = \frac{kb_0}{2}, \quad a_1 = \frac{k}{2}(b_1 - b_0), \quad a_2 = \frac{-kb_1}{2}, \quad k = \pm 2\sqrt{\beta + \alpha^2}. \quad (31)$$

Substituting (31) into (14) with (5), (10) and (12), we obtain the following solutions of Eq. (9):

$$\psi_{5,6}(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{1 + \kappa e^{\pm 2\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0}} \left(1 - \frac{1}{1 + \kappa e^{\pm 2\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0}} \right) \right\}^{1/2} e^{i(\alpha x + \beta t)}, \quad (32)$$

$$\psi_{7,8}(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{1 + \kappa e^{\pm 2\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0}} \left(1 - \frac{1}{1 + \kappa e^{\pm 2\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0}} \right) \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (33)$$

If we set $\kappa = 1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \tanh(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}, \quad (34)$$

$$\psi(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \tanh(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (35)$$

If we set $\kappa = -1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}, \quad (36)$$

$$\psi(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (37)$$

Result 3.

$$a_0 = \frac{3kb_0}{4}, \quad a_1 = -2kb_0, \quad a_2 = kb_0, \quad b_1 = -2b_0, \quad k = \pm 2\sqrt{\frac{2(\beta + \alpha^2)}{5}}. \quad (38)$$

Substituting (38) into (14) with (5), (10) and (12), we obtain the following solutions of Eq. (9):

$$\psi_{9,10}(x, t) = \pm \left\{ \frac{1}{2} \sqrt{\frac{2(\beta + \alpha^2)}{5}} \left(3 - \frac{2}{1 + \kappa e^{\pm 2\sqrt{2(\beta + \alpha^2)/5}(x - 2\alpha t) + \xi_0}} \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (39)$$

$$\psi_{11,12}(x, t) = \pm i \left\{ \frac{1}{2} \sqrt{\frac{2(\beta + \alpha^2)}{5}} \left(3 - \frac{2}{1 + \kappa e^{\pm 2\sqrt{2(\beta + \alpha^2)/5}(x - 2\alpha t) + \xi_0}} \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}. \quad (40)$$

If we set $\kappa = 1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{1}{2} \sqrt{\frac{2(\beta + \alpha^2)}{5}} \left(2 + \tanh \left(\pm \sqrt{\frac{2(\beta + \alpha^2)}{5}}(x - 2\alpha t) + \xi_0 \right) \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (41)$$

$$\psi(x, t) = \pm i \left\{ \frac{1}{2} \sqrt{\frac{2(\beta + \alpha^2)}{5}} \left(2 + \tanh \left(\pm \sqrt{\frac{2(\beta + \alpha^2)}{5}}(x - 2\alpha t) + \xi_0 \right) \right) \right\}^{1/2} e^{i(\alpha x + \beta t)}. \quad (42)$$

If we set $\kappa = -1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{1}{2} \sqrt{\frac{2(\beta + \alpha^2)}{5}} \left(2 + \coth \left(\pm \sqrt{\frac{2(\beta + \alpha^2)}{5}}(x - 2\alpha t) + \xi_0 \right) \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (43)$$

$$\psi(x, t) = \pm i \left\{ \frac{1}{2} \sqrt{\frac{2(\beta + \alpha^2)}{5}} \left(2 + \coth \left(\pm \sqrt{\frac{2(\beta + \alpha^2)}{5}}(x - 2\alpha t) + \xi_0 \right) \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}. \quad (44)$$

Result 4.

$$a_0 = 0, \quad a_1 = 0, \quad a_2 = kb_0, \quad b_1 = -2b_0, \quad k = \pm\sqrt{\beta + \alpha^2}. \quad (45)$$

Substituting (45) into (14) with (5), (10) and (12), we obtain the following solutions of Eq. (9):

$$\psi_{13,14}(x, t) = \pm \left\{ \sqrt{\beta + \alpha^2} \left(\frac{(1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-2}}{1 - 2(1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-1}} \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (46)$$

$$\psi_{15,16}(x, t) = \pm i \left\{ \sqrt{\beta + \alpha^2} \left(\frac{(1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-2}}{1 - 2(1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-1}} \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}. \quad (47)$$

If we set $\kappa = \pm 1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (48)$$

$$\psi(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}. \quad (49)$$

Result 5.

$$a_0 = kb_0, \quad a_1 = -2kb_0, \quad a_2 = kb_0, \quad b_1 = -2b_0, \quad k = \pm\sqrt{\beta + \alpha^2}. \quad (50)$$

Substituting (50) into (14) with (5), (10) and (12), we obtain the following solutions of Eq. (9):

$$\psi_{17,18}(x, t) = \pm \left\{ \sqrt{\beta + \alpha^2} \left(\frac{((1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-1} - 1)^2}{1 - 2(1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-1}} \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (51)$$

$$\psi_{19,20}(x, t) = \pm i \left\{ \sqrt{\beta + \alpha^2} \left(\frac{((1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-1} - 1)^2}{1 - 2(1 + \kappa e^{\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0})^{-1}} \right) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}. \quad (52)$$

If we set $\kappa = \pm 1$, we obtain

$$\psi(x, t) = \pm \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \coth(\pm\sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}, \quad (53)$$

$$\psi(x, t) = \pm i \left\{ \frac{\sqrt{\beta + \alpha^2}}{2} (1 + \coth(\pm \sqrt{\beta + \alpha^2}(x - 2\alpha t) + \xi_0)) \right\}^{1/2} \times e^{i(\alpha x + \beta t)}. \quad (54)$$

4 Conclusions

In this paper, we have proposed the generalized Kudryashov method for solving the Enkhaus equation. This work has illustrated that the solutions obtained in [30] are considered as a special case of our obtained solutions and a new results have been obtained using this method. This method is direct, effective and can be extended for solving many systems of nonlinear PDEs. The soliton solutions that are retrievable from this equation are topological and singular soliton solutions only. Being a dissipative model, it is not possible to obtain non-topological soliton solution. Therefore, it makes sense that this model only retrieves singular and topological soliton solutions.

Acknowledgment. The authors express their sincere thanks to the referees for their careful review of this manuscript and their useful suggestions.

References

1. M.A. Abdou, The extended F -expansion method and its application for a class of nonlinear evolution equations, *Chaos Solitons Fractals*, **31**:95–104, 2007.
2. A. Biswas, Temporal 1-soliton solution of the complex Ginzburg–Landau equation with power law nonlinearity, *Progress in Electromagnetics Research*, **96**:1–7, 2009.
3. A. Biswas, 1-soliton solution of the $K(m, n)$ equation with generalized evolution and time-dependent damping and dispersion, *Comput. Math. Appl.*, **59**(8):2538–2542, 2010.
4. A. Biswas, K. Khan, A. Rahman, A. Yildirim, T. Hayat, O.M. Aldossary, Bright and dark optical solitons in birefringent fibers with Hamiltonian perturbations and Kerr law nonlinearity, *J. Optoelectron. Adv. Mater.*, **14**(7):571–576, 2012.
5. A. Biswas, D. Milovic, A. Ranasinghe, Solitary waves of Boussinesq equation in a power law media, *Commun. Nonlinear Sci. Numer. Simul.*, **14**(11):3738–3742, 2009.
6. A. Biswas, D. Milovic, M. Savescu, M.F. Mahmood, K.R. Khan, R. Kohl, Optical solitons perturbation in nanofibers with improved nonlinear Schrödinger's equation by semi-inverse variational principle, *J. Nonlinear Opt. Phys. Mater.*, **12**(4):1250054, 2012.
7. A. Biswas, C. Zony, E. Zerrad, Soliton perturbation theory for the quadratic nonlinear Klein–Gordon equation, *Appl. Math. Comput.*, **203**(1):153–156, 2008.
8. H. Bulut, Y. Pandir, S.T. Demiray, Exact solutions of time-fractional KdV equations by using generalized Kudryashov method, *International Journal of Modeling and Optimization*, **4**(4):315–320, 2014.
9. F. Calogero, W. Eckhaus, Nonlinear evolution equations, rescalings, model PDEs and their integrability, *Inverse Probl.*, **3**:229–262, 1987.

10. S.A. EL-Wakil, M.A. Abdou, New exact travelling wave solutions using modified extended tanh-function method, *Chaos Solitons Fractals*, **31**:840–852, 2007.
11. S.D. Lillo F. Calogero, The Eckhaus PDE $i\psi_t + \psi_{xx} + 2(|\psi|^2)_x\psi + |\psi|^4\psi = 0$, *Inverse Probl.*, **3**:633–681, 1987.
12. A.L. Fabian, R. Kohl, A. Biswas, Perturbation of topological solitons due to sine-Gordon equation and its type, *Commun. Nonlinear Sci. Numer. Simul.*, **14**(4):1227–1244, 2009.
13. E. Fan, J. Zhang, Applications of the Jacobi elliptic function method to special-type nonlinear equations, *Phys. Lett. A*, **305**:383–392, 2002.
14. J.H. He, X.H. Wu, Exp-function method for nonlinear wave equations, *Chaos Solitons Fractals*, **30**:700–708, 2006.
15. K.R. Khan, M.F. Mahmood, A. Biswas, Coherent super continuum generation in photonic crystal fibers at visible and near infrared wavelengths, *IEEE J. Sel. Top. Quantum Electron.*, **20**(5):7500309, 2014.
16. N.A. Kudryashov, Exact soliton solutions of the generalized evolution equation of wave dynamics, *J. Appl. Math. Mech.*, **52**:361–365, 1988.
17. N.A. Kudryashov, Exact solutions of the generalized Kuramoto–Sivashinsky equation, *Phys. Lett. A*, **147**:287–291, 1990.
18. N.A. Kudryashov, On types of nonlinear nonintegrable equations with exact solutions, *Phys. Lett. A*, **155**:269–275, 1991.
19. N.A. Kudryashov, One method for finding exact solutions of nonlinear differential equations, *Commun. Nonlinear Sci. Numer. Simul.*, **17**:2248–2253, 2012.
20. N.A. Kudryashov, A.S. Zakharchenko, A note on solutions of the generalized Fisher equation, *Appl. Math. Lett.*, **32**:53–56, 2014.
21. M. Mirzazadeh, M. Eslami, A. Biswas, Optical solitons by Kudrashov’s method, *Optik*, **125**(23):6874–6880, 2014.
22. P.N. Ryabov, Exact solutions of the Kudryashov–Sinelschchikov equation, *Appl. Math. Comput.*, **217**:3585–3590, 2010.
23. M. Savescu, K.R. Khan, R.W. Kohl, L. Moraru, A. Yildirim, A. Biswas, Optical soliton perturbation with improved nonlineaer Schrödinger’s equation in nanofibers, *J. Nanoelectron. Optoelectron.*, **8**(2):208–220, 2013.
24. M. Savescu, K.R. Khan, P. Naruka, H. Jafari, L. Moraru, A. Biswas, Optical solitons in photonic nanowaveguides with an improved nonlinear Schrödinger’s equation, *J. Comput. Theor. Nanosci.*, **10**(5):1182–1191, 2013.
25. H. Triki, S. Crutcher, A. Yildirim, T. Hayat, O.M. Aldossary, A. Biswas, Bright and dark solitons of the modified complex Ginzburg–Landau equation with parabolic and dual-power law nonlinearity, *Romanian Reports in Physics*, **64**(2):367–370, 2012.
26. M.L. Wang, J.L. Zhang, X.Z. Li, The (G'/G) -expansion method and traveling wave solutions of nonlinear evolutions equations in mathematical physics, *Phys. Lett. A*, **372**:417–423, 2008.
27. A.M. Wazwaz, A sine-cosine method for handling nonlinear wave equations, *Math. Comput. Modelling*, **40**:499–508, 2004.
28. A.M. Wazwaz, The tanh method for travelling wave solutions of nonlinear equations, *Appl. Math. Comput.*, **154**:714–723, 2004.

29. A.M. Wazwaz, The extended tanh method for abundant solitary wave solutions of nonlinear wave equations, *Appl. Math. Comput.*, **187**:1131–1142, 2007.
30. H. Zhang, New application of the (G'/G) -expansion method, *Commun. Nonlinear Sci. Numer. Simul.*, **14**:3220–3225, 2009.
31. J.L. Zhang, M.L. Wang, Y.M. Wang, Z.D. Fang, The improved F -expansion method and its applications, *Phys. Lett. A*, **350**:103–109, 2006.
32. X.Q. Zhao, H.Y. Zhi, H.Q. Zhang, Improved Jacobi elliptic function method with symbolic computation to construct new double-periodic solutions for the generalized Ito system, *Chaos Solitons Fractals*, **28**:112–126, 2006.
33. Q. Zhou, Q. Zhu, A. Biswas, Optical solitons in birefringent fibers with parabolic law nonlinearity, *Opt. Appl.*, **41**(3):399–409, 2014.
34. Q. Zhou, Q. Zhu, Y. Liu, P. Yao, A.H. Bhrawy, L. Moraru, A. Biswas, Brightdark combo optical solitons with non-local nonlinearity in parabolic law medium, *Optoelectron. Adv. Mater. Rapid Commun.*, **8**(9):837–839, 2014.
35. Q. Zhou, Q. Zhu, Y. Liu, H. Yu, P. Yao, A. Biswas, Thirring optical solitons in birefringent fibers with spatio-temporal dispersion and Kerr law nonlinearity, *Laser Phys.*, **25**(1):015402, 2015.
36. Q. Zhou, Q. Zhu, L. Moraru, A. Biswas, Optical solitons with spatially dependent coefficients by Lie symmetry, *Optoelectron. Adv. Mater. Rapid Commun.*, **8**(7):800–803, 2014.