

# On the influence of the arithmetical character of the parameters for the Lerch zeta-function

Jolita IGNATAVIČIŪTĖ (VU)

*e-mail: jolita.ignataviciute@maf.vu.lt*

## 1. Introduction

Let  $s = \sigma + it$  with  $\sigma, t \in \mathbb{R}$ ,  $i^2 = -1$ , be a complex variable. In 1887 M. Lerch introduced the function, defined on the half-plane  $\sigma > 1$  by the following absolutely convergent Dirichlet series

$$L(\lambda, \alpha, s) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda m}}{(m + \alpha)^s}.$$

The function  $L(\lambda, \alpha, s)$  depends on two fixed parameters  $\lambda \in \mathbb{R}$  and  $\alpha \in \mathbb{R}_+$ . Without loss of generality we consider  $0 < \alpha \leq 1$ . The investigations of this function in different aspects show that its properties are strongly effected by the arithmetical character of the parameters  $\lambda$  and  $\alpha$ . In this paper we restrict ourselves to the discrete value-distribution of  $L(\lambda, \alpha, s)$ .

Let  $h \in \mathbb{R}_+$  be a fixed number, and  $k, N \in \mathbb{N} \cup \{0\}$ . We consider probability measures of the following form

$$\mu_N(\dots) = \frac{1}{N+1} \# \{k \in [0, N]: \dots\},$$

where in place of dots some condition satisfied  $k$  is to be written. Let  $\mathcal{B}(S)$  stand for the class of Borel sets of the space  $S$ . Let  $D = \{s \in \mathbb{C}: \sigma > 1/2\}$ . We write  $H(D)$  for the space of analytic on  $D$  functions equipped with the topology of uniform convergence on compacta. The notation  $M(D)$  means the space of meromorphic on  $D$  functions equipped with the topology as was stated above. Consider the following three probability measures:

$$\begin{aligned} P_{1N}(A) &= \mu_N(L(\lambda, \alpha, \sigma + ikh) \in A), & A \in \mathcal{B}(\mathbb{C}), & \lambda \notin \mathbb{Z}, \\ P_{2N}(A) &= \mu_N(L(\lambda, \alpha, s + ikh) \in A), & A \in \mathcal{B}(H(D)), & \lambda \notin \mathbb{Z}, \\ P_{3N}(A) &= \mu_N(L(\lambda, \alpha, s + ikh) \in A), & A \in \mathcal{B}(M(D)), & \lambda \in \mathbb{Z}. \end{aligned}$$

In recent years the discrete value-distribution for the Lerch zeta-function with a transcendental parameter  $\alpha$  was studied by the author. The main results are as follows.

Define  $\gamma$  to be the unit circle on  $\mathbb{C}$ , and

$$\Omega_1 = \prod_{m=0}^{\infty} \gamma_m,$$

where  $\gamma_m = \gamma$ ,  $m = 0, 1, 2, \dots$ . There exists the probability Haar measure  $m_{1H}$  on  $(\Omega_1, \mathcal{B}(\Omega_1))$ . Let  $\omega_1(m)$  be the projection of  $\omega_1 \in \Omega_1$  to  $\gamma_m$ . Define the  $\mathbb{C}$ -valued random element on  $(\Omega_1, \mathcal{B}(\Omega_1), m_{1H})$  by

$$L(\lambda, \alpha, \sigma, \omega_1) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda m} \omega_1(m)}{(m + \alpha)^\sigma}, \quad \sigma \in D, \quad \omega_1 \in \Omega_1.$$

**Theorem 1.** *Suppose that  $\lambda \notin \mathbb{Z}$ ,  $\alpha$  is a transcendental number, and  $\exp\{2\pi/h\}$  is a rational number. Then the probability measure  $P_{1N}$  converges weakly to the distribution of  $L(\lambda, \alpha, \sigma, \omega_1)$  as  $N \rightarrow \infty$ .*

For the proof see [2].

We obtain an  $H(D)$ -valued random element on  $(\Omega_1, \mathcal{B}(\Omega_1), m_{1H})$  for  $s$  in place of  $\sigma$  in  $L(\lambda, \alpha, \sigma, \omega_1)$ .

**Theorem 2.** *Under the assumptions of Theorem 1 the probability measure  $P_{2N}$  converges weakly to the distribution of  $L(\lambda, \alpha, s, \omega_1)$  as  $N \rightarrow \infty$ .*

For the proof see [1] and the references given there.

**Theorem 3.** *Suppose that  $\lambda \in \mathbb{Z}$ ,  $\alpha$  is a transcendental number, and  $\exp\{2\pi/h\}$  is a rational number. Then the probability measure  $P_{3N}$  converges weakly to the distribution of  $L(\lambda, \alpha, s, \omega_1)$  as  $N \rightarrow \infty$ .*

See [3] for more details.

For a rational parameter  $\alpha$  the results are as follows.

**Theorem 1'.** *Suppose that  $\lambda \notin \mathbb{Z}$ ,  $\alpha \in \mathbb{Q}$ , and  $\exp\{2\pi k/h\}$ ,  $k \in \mathbb{Z}$ ,  $k \neq 0$ , is an irrational number. Then there exists a probability measure  $P_1$  on  $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$  such that  $P_{1N}$  converges weakly to  $P_1$  as  $N \rightarrow \infty$ .*

**Theorem 2'.** *Under the assumptions of Theorem 1' there exists a probability measure  $P_2$  on  $(H(D), \mathcal{B}(H(D)))$  such that  $P_{2N}$  converges weakly to  $P_2$  as  $N \rightarrow \infty$ .*

**Theorem 3'.** *Suppose that  $\lambda \in \mathbb{Z}$ ,  $\alpha \in \mathbb{Q}$ , and  $\exp\{2\pi k/h\}$ ,  $k \in \mathbb{Z}$ ,  $k \neq 0$ , is an irrational number. Then there exists a probability measure  $P_3$  on  $(H(D), \mathcal{B}(H(D)))$  such that  $P_{3N}$  converges weakly to  $P_3$  as  $N \rightarrow \infty$ .*

The explicit form of the limit measure is obtained for  $\alpha = a/b$ ,  $1 \leq a \leq b$ ,  $(a, b) = 1$ .  
Set

$$\Omega_2 = \prod_p \gamma_p,$$

where  $\gamma_p = \gamma$  for all prime numbers  $p$ . There exists the probability Haar measure  $m_{2H}$  on  $(\Omega_2, \mathcal{B}(\Omega_2))$ . Let  $\omega_2(p)$  stand for the projection of  $\omega_2 \in \Omega_2$  to  $\gamma_p$ . For  $m \in \mathbb{N} \cup \{0\}$ , set

$$\omega(m) = \prod_{p^\alpha \parallel m} \omega^\alpha(p), \quad \alpha \in \mathbb{N}.$$

Define the  $\mathbb{C}$ -valued random element on  $(\Omega_2, \mathcal{B}(\Omega_2), m_{2H})$  by

$$L(\lambda, \alpha, \sigma, \omega_2) = \omega_2(b) b^\sigma e^{-2\pi i \lambda a/b} \sum_{\substack{m=1 \\ m \equiv a \pmod{b}}}^{\infty} \frac{e^{2\pi i \lambda m/b} \omega_2(m)}{m^\sigma}, \quad \sigma \in D, \quad \omega_2 \in \Omega_2.$$

**Theorem 1''.** *Suppose that  $\lambda \notin \mathbb{Z}$ ,  $\mathbb{Q} \ni \alpha = a/b$ ,  $1 \leq a \leq b$ ,  $(a, b) = 1$ , and  $\exp\{2\pi k/h\}$ ,  $k \in \mathbb{Z}$ ,  $k \neq 0$ , is an irrational number. Then the probability measure  $P_{1N}$  converges weakly to the distribution of  $L(\lambda, \alpha, \sigma, \omega_2)$  as  $N \rightarrow \infty$ .*

It follows easily that for  $s$  in place of  $\sigma$  in  $L(\lambda, \alpha, \sigma, \omega_2)$  we obtain an  $H(D)$ -valued random element  $L(\lambda, \alpha, s, \omega_2)$  on  $(\Omega_2, \mathcal{B}(\Omega_2), m_{2H})$ .

**Theorem 2''.** *Under the assumptions of Theorem 1'' the probability measure  $P_{2N}$  converges weakly to the distribution of  $L(\lambda, \alpha, s, \omega_2)$  as  $N \rightarrow \infty$ .*

**Theorem 3''.** *Suppose that  $\lambda \in \mathbb{Z}$ ,  $\mathbb{Q} \ni \alpha = a/b$ ,  $1 \leq a \leq b$ ,  $(a, b) = 1$ , and  $\exp\{2\pi k/h\}$ ,  $k \in \mathbb{Z}$ ,  $k \neq 0$ , is an irrational number. Then the probability measure  $P_{3N}$  converges weakly to the distribution of  $L(\lambda, \alpha, s, \omega_2)$  as  $N \rightarrow \infty$ .*

In this article we present the proof of Theorem 3''. The other results in the case of a rational parameter  $\alpha$  may be proved in much the same way.

## 2. Proof of Theorem 3''

For  $\lambda \in \mathbb{Z}$  the function  $L(\lambda, \alpha, s)$  reduces to the Hurwitz zeta-function

$$\zeta(\alpha, s) = \sum_{m=0}^{\infty} \frac{1}{(m + \alpha)^s}.$$

This function is analytically continuable over the whole complex plane except for a simple pole at the point  $s = 1$  with the residue 1. We may interpret the function  $\zeta(\alpha, s)$  as

$$\zeta(\alpha, s) = \frac{f_2(\alpha, s)}{f_1(s)},$$

where

$$\begin{aligned} f_1(s) &= 1 - 2^{1-s}, \\ f_2(\alpha, s) &= \zeta(\alpha, s)f_1(s). \end{aligned}$$

Moreover, for  $\mathbb{Q} \ni \alpha = a/b, 1 \leq a \leq b, (a, b) = 1$ , the function  $f_2(\alpha, s)$  can be interpreted as a product of entire functions, i.e.,

$$\begin{aligned} f_2(\alpha, s) &= f_1(s) \sum_{m=0}^{\infty} \frac{1}{(m + \alpha)^s} = f_1(s) \sum_{m=0}^{\infty} \frac{1}{(m + a/b)^s} \\ &= f_1(s)b^s \sum_{m=0}^{\infty} \frac{1}{(mb + a)^s} = f_1(s)b^s \sum_{\substack{m=1 \\ m \equiv a \pmod{b}}}^{\infty} \frac{1}{m^s} \stackrel{def}{=} f_1(s)g_1(s)g_2(s), \end{aligned}$$

where

$$g_1(s) = b^s, \quad g_2(s) = \sum_{\substack{m=1 \\ m \equiv a \pmod{b}}}^{\infty} \frac{1}{m^s}.$$

By the same method as in [1] it follows that the probability measures

$$\mu_N(f_1(s + ikh) \in A), \quad \mu_N(g_i(s + ikh) \in A), \quad A \in \mathcal{B}(H(D)), \quad i = 1, 2,$$

converge weakly to  $P_{f_1}, P_{g_i}, i = 1, 2$ , respectively, as  $N \rightarrow \infty$ , where

$$\begin{aligned} P_{f_1}(A) &= m_{2H}(\omega_2 \in \Omega_2: f_1(s, \omega_2) \in A), \\ P_{g_i}(A) &= m_{2H}(\omega_2 \in \Omega_2: g_i(s, \omega_2) \in A), \end{aligned}$$

$i = 1, 2, s \in D, A \in \mathcal{B}(H(D))$ , and

$$\begin{aligned} f_1(s, \omega_2) &= (1 - 2^{1-s})\omega_2(2), \\ g_1(s, \omega_2) &= b^s\omega_2(b), \quad g_2(s, \omega_2) = \sum_{\substack{m=1 \\ m \equiv a \pmod{b}}}^{\infty} \frac{\omega_2(m)}{m^s}. \end{aligned}$$

The rest of the proof runs as in [3], p. 16–20, with the auxiliary function  $u: H(D) \times H(D) \times H(D) \rightarrow H(D)$  defined by the formulae

$$u(h_1, h_2, h_3) = h_1 * h_2 * h_3, \quad h_1, h_2, h_3 \in H(D).$$

We obtain that

$$\begin{aligned} \mu_N(f_2(\alpha, s + ikh) \in A) \\ = \mu_N(f_1(s + ikh)g_1(s + ikh)g_2(s + ikh) \in A), \quad A \in \mathcal{B}(H(D)), \end{aligned}$$

converges weakly to

$$P_{f_2}(A) = m_{2H}(\omega_2 \in \Omega_2: f_1(s, \omega_2)g_1(s, \omega_2)g_2(s, \omega_2) \in A),$$

$s \in D, \omega_2 \in \Omega_2, A \in \mathcal{B}(H(D))$ , as  $N \rightarrow \infty$ .

Finally, applying the auxiliary function  $v: H(D) \times H(D) \rightarrow M(D)$  defined by the formulae

$$v(h_1, h_2) = \frac{h_1}{h_2}, \quad h_1, h_2 \in H(D),$$

we obtain the assertion of the theorem.

## References

- [1] J. Ignatavičiūtė, A limit theorem for the Lerch zeta-function, *Liet. Matem. Rink.*, **40** (special issue), 21–27 (2000).
- [2] J. Ignatavičiūtė, On statistic properties of the Lerch zeta-function, *Liet. Matem. Rink.*, **41**(4), 424–440 (2001) (in Russian).
- [3] J. Ignatavičiūtė, On statistic properties of the Lerch zeta-function, II. *Liet. Matem. Rink.*, (to appear(2002)).

## Parametų aritmetinės prigimties įtaka Lercho dzeta funkcijai

J. Ignatavičiūtė

Įrodoma diskreti ribinė teorema Lercho dzeta funkcijai su parametrais  $\alpha = a/b, a \in \mathbb{Z}, b \in \mathbb{N}, (a, b) = 1$ , ir  $\lambda \in \mathbb{Z}$  meromorfinių pusplokštumėje  $\sigma > 1/2$  funkcijų erdvėje. Pateikiami analogiški rezultatai kompleksinėje plokštumoje bei analizinė pusplokštumėje  $\sigma > 1/2$  funkcijų erdvėje.