

Impulsive coupled systems with generalized jump conditions

Feliz Minhós^a, Robert de Sousa^b

^aDepartamento de Matemática, Escola de Ciências e Tecnologia,
Centro de Investigação em Matemática e Aplicações (CIMA),
Instituto de Investigação e Formação Avançada,
Universidade de Évora,
Rua Romão Ramalho, 59, 7000-671 Évora, Portugal
fminhos@uevora.pt

^bFaculdade de Ciências e Tecnologia,
Núcleo de Matemática e Aplicações (NUMAT),
Universidade de Cabo Verde,
Campus de Palmarejo, 279 Praia, Cabo Verde
robert.sousa@docente.unicv.edu.cv

Received: May 24, 2017 / **Revised:** October 3, 2017 / **Published online:** December 14, 2017

Abstract. This work considers a second-order impulsive coupled system with full nonlinearities, generalized impulse functions and mixed boundary conditions. This is the first time where such coupled systems are considered with nonlinearities with dependence on both unknown functions and their derivatives, together impulsive functions given by more general framework allowing jumps on the both functions and both derivatives.

The arguments apply the fixed point theory, Green's functions technique, L^1 -Carathéodory functions theory and Schauder's fixed point theorem.

An application to the transverse vibration system of elastically coupled double-string is presented in the last section.

Keywords: impulsive coupled systems, L^1 -Carathéodory functions, Green's functions, Schauder's fixed point theorem, elastically coupled double-string system.

1 Introduction

In this paper, we consider the second-order impulsive coupled system with mixed boundary conditions

$$\begin{aligned}u''(x) &= f(x, u(x), u'(x), v(x), v'(x)), \\v''(x) &= h(x, u(x), u'(x), v(x), v'(x)), \\u(a) &= A_1, \quad u'(b) = B_1, \\v(a) &= A_2, \quad v(b) = B_2,\end{aligned}\tag{1}$$

where $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ are L^1 -Carathéodory functions, $A_1, A_2, B_1, B_2 \in \mathbb{R}$, with the generalized impulsive conditions

$$\begin{aligned}\Delta u(x_k) &= I_{0k}(x_k, u(x_k), u'(x_k)), \\ \Delta u'(x_k) &= I_{1k}(x_k, u(x_k), u'(x_k)), \quad k = 1, 2, \dots, n, \\ \Delta v(\tau_j) &= J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)), \\ \Delta v'(\tau_j) &= J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)), \quad j = 1, 2, \dots, m,\end{aligned}\tag{2}$$

where, for $i = 0, 1$, $\Delta u^{(i)}(x_k) = u^{(i)}(x_k^+) - u^{(i)}(x_k^-)$, $\Delta v^{(i)}(\tau_j) = v^{(i)}(\tau_j^+) - v^{(i)}(\tau_j^-)$, and being $I_{ik}, J_{ij} \in C([a, b] \times \mathbb{R}^2, \mathbb{R})$ with x_k, τ_j fixed points such that $a < x_1 < x_2 < \dots < x_n < b$ and $a < \tau_1 < \tau_2 < \dots < \tau_m < b$.

The theory of impulsive differential equations describes processes in which a sudden change of state occurs at certain moments. Several authors (see, for example, [1, 3, 6–8, 10–14, 17, 20, 25]) have dealt with impulsive differential equations from different points of view and using many techniques.

There are many phenomena and applications related to impulsive differential systems, for example, we can find biological models, population dynamics, neural networks, models in economics, on time scales, on state-dependent delays, on delay-dependent impulsive control, on electrochemical communication between cells in the brain (see for instance, [2, 4, 5, 9, 15, 16, 19, 22–24, 26, 27, 29, 30]), among others.

In [28], the author considers a sufficient conditions for the existence and uniqueness of solutions to the following complex dynamical network in the form of a coupled system of $m + 2$ point boundary conditions for impulsive fractional differential equations

$$\begin{aligned}{}^c D^\alpha u(t) &= \phi(t, u(t), v(t)), \quad t \in [0, 1], \quad t \neq t_j, \quad j = 1, \dots, m, \\ {}^c D^\beta v(t) &= \psi(t, u(t), v(t)), \quad t \in [0, 1], \quad t \neq t_i, \quad i = 1, \dots, n, \\ u(0) &= h(u), \quad u(1) = g(u) \quad \text{and} \quad v(0) = k(v), \quad v(1) = f(v), \\ \Delta u(t_j) &= I_j(u(t_j)), \quad \Delta u'(t_j) = \bar{I}_j(u(t_j)), \quad j = 1, \dots, m, \\ \Delta v(t_i) &= I_i(v(t_i)), \quad \Delta v'(t_i) = \bar{I}_i(v(t_i)), \quad i = 1, \dots, n,\end{aligned}$$

where $1 < \alpha, \beta \leq 2$, $\phi, \psi : [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ are continuous functions, and $g, h : X \rightarrow \mathbb{R}$, $f, k : Y \rightarrow \mathbb{R}$ are continuous functionals defined by

$$\begin{aligned}g(u) &= \sum_{j=1}^p \lambda_j u(\xi_j), & h(u) &= \sum_{j=1}^p \lambda_j u(\eta_j), \\ f(v) &= \sum_{i=1}^q \delta_i v(\xi_i), & k(v) &= \sum_{i=1}^q \delta_i v(\eta_i),\end{aligned}$$

$\xi_i, \eta_i, \xi_j, \eta_j \in (0, 1)$ for $i = 1, \dots, q$ and $j = 1, \dots, p$.

In [18], it is studied the BVP for second-order singular differential system on the whole line with impulse effects, i.e., consisting of the differential system

$$\begin{aligned} [\phi_p(\rho(t)x'(t))] &= f(t, x(t), y(t)), \quad \text{a.e. } t \in \mathbb{R}, \\ [\phi_q(\varrho(t)y'(t))] &= g(t, x(t), y(t)), \quad \text{a.e. } t \in \mathbb{R}, \end{aligned}$$

subjected to the boundary conditions

$$\lim_{t \rightarrow \pm\infty} x(s) = 0, \quad \lim_{t \rightarrow \pm\infty} y(s) = 0$$

and the impulse effects

$$\begin{aligned} \Delta x(t_k) &= I_k(t_k, x(t_k), y(t_k)), \quad k \in \mathbb{Z}, \\ \Delta y(t_k) &= J_k(t_k, x(t_k), y(t_k)), \quad k \in \mathbb{Z}, \end{aligned}$$

where

- (i) $\rho, \varrho \in C^0(\mathbb{R}, [0, \infty))$, $\rho(t), \varrho(t) > 0$ for all $t \in \mathbb{R}$ with $\int_{-\infty}^{+\infty} ds/\rho(s) < +\infty$ and $\int_{-\infty}^{+\infty} ds/\varrho(s) < +\infty$,
- (ii) $\phi_p(x) = |x|^{p-2}x$, $\phi_q(x) = |x|^{q-2}x$ with $p > 1$ and $q > 1$ are Laplace operators,
- (iii) f, g on \mathbb{R}^3 are Carathéodory functions,
- (iv) $\dots < t_k < t_{k+1} < t_{k+2} < \dots$ with $\lim_{k \rightarrow -\infty} t_k = -\infty$ and $\lim_{k \rightarrow +\infty} t_k = +\infty$, $\Delta x(t_k) = u(t_k^+) - x(t_k^-)$ and $\Delta y(t_k) = y(t_k^+) - y(t_k^-)$ ($k \in \mathbb{Z}$), \mathbb{Z} is the set of all integers,
- (v) $\{I_k\}, \{J_k\}$, with $I_k, J_k : \mathbb{R}^3 \rightarrow \mathbb{R}$, are Carathéodory sequences.

Motivated by these works, we follow arguments applied in [21] to study problem (1)–(2). We point out that is the first time when second-order coupled differential equations systems include full nonlinearities. That is, they depend on the unknown functions, and their first derivatives, together with generalized impulsive conditions with dependence on the first derivative, too.

The paper is organized as it follows: Section 2 contains the preliminary results: definitions and some auxiliary lemmas. Section 3 contains the main result: an existence solution of the problem. In the last section, our main theorem is illustrated by an example and applications to a real phenomena: the transverse vibrations system of elastically coupled double-string model.

2 Definitions and auxiliary results

Define $u(x_k^\pm) := \lim_{x \rightarrow x_k^\pm} u(x)$, consider the set

$$PC_1([a, b]) = \left\{ u: [a, b] \rightarrow \mathbb{R}, u \text{ is continuous for } x \neq x_k, u(x_k) = u(x_k^-), u(x_k^+) \text{ exists for } k = 1, 2, \dots, n \right\}$$

and the space $X_1 := PC_1^1([a, b]) = \{u: u'(t) \in PC_1([a, b])\}$ equipped with the norm $\|u\|_{X_1} = \max\{\|u\|, \|u'\|\}$, where

$$\|w\| := \sup_{x \in [a, b]} |w(x)|.$$

Analogously, define the space $X_2 := PC_2^1([a, b]) = \{v: v'(x) \in PC_2([a, b])\}$, where

$$PC_2([a, b]) = \{v: v[a, b] \rightarrow \mathbb{R}, v \text{ is continuous for } \tau \neq \tau_j, v(\tau_j) = v(\tau_j^- v), \\ v(\tau_j^+) \text{ exists for } j = 1, 2, \dots, m\},$$

equipped with the norm $\|v\|_{X_2} = \max\{\|v\|, \|v'\|\}$.

Denoting $X := X_1 \times X_2$ and the norm $\|(u, v)\|_X = \max\{\|u\|_{X_1}, \|v\|_{X_2}\}$, it is clear that $(X, \|\cdot\|_X)$ is a Banach space.

A pair of functions (u, v) is a solution of problem (1)–(2) if $(u, v) \in X$ and verifies conditions (1) and (2).

Definition 1. A function $g : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is L^1 -Carathéodory if

- (i) for each $(t, y, z, w) \in \mathbb{R}^4$, $x \mapsto g(x, t, y, z, w)$ is measurable on $[a, b]$;
- (ii) for a.e. $x \in [a, b]$, $(t, y, z, w) \mapsto g(x, t, y, z, w)$ is continuous on \mathbb{R}^4 ;
- (iii) for each $\rho > 0$, there exists a positive function $\phi_\rho \in L^1([a, b])$, and for $(t, y, z, w) \in \mathbb{R}^4$ such that

$$\max\{|t|, |y|, |z|, |w|\} < \rho,$$

one has

$$|g(x, t, y, z, w)| \leq \phi_\rho(t), \quad \text{a.e. } x \in [a, b].$$

Lemma 1. A pair of functions $(u, v) \in X$ is a solution of problem (1)–(2) if and only if

$$u(x) = A_1 + B_1(x - a) \\ + \sum_{x_k < x} [I_{0k}(x_k, u(x_k), u'(x_k)) + I_{1k}(x_k, u(x_k), u'(x_k))(x - x_k)] \\ - (x - a) \sum_{k=1}^n I_{1k}(x_k, u(x_k), u'(x_k)) \\ + \int_a^b G_1(x, s) f(s, u(s), u'(s), v(s), v'(s)) ds$$

with $G_1(x, s)$ given by

$$G_1(x, s) = \begin{cases} a - s, & a \leq x \leq s \leq b, \\ a - x, & a \leq s \leq x \leq b, \end{cases} \quad (3)$$

and

$$\begin{aligned}
 v(x) = & A_2 + \frac{B_2 - A_2}{b - a}(x - a) \\
 & + \sum_{\tau_j < x} [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j)] \\
 & - \frac{x - a}{b - a} \sum_{j=1}^m [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j)] \\
 & + \int_a^b G_2(x, s)h(s, u(s), u'(s), v(s), v'(s)) ds
 \end{aligned}$$

with $G_2(x, s)$ defined by

$$G_2(x, s) = \frac{1}{a - b} \begin{cases} (a - s)(b - x), & a \leq x \leq s \leq b, \\ (x - a)(b - s), & a \leq s \leq x \leq b. \end{cases} \tag{4}$$

The proof follows standard calculus and it is omitted.

A key tool is the Schauder’s fixed point theorem:

Theorem 1. (See [31].) *Let Y be a nonempty, closed, bounded, and convex subset of a Banach space X , and suppose that $P : Y \rightarrow Y$ is a compact operator. Then P has at least one fixed point in Y .*

3 Main theorem

The main result will provide the existence of at least one solution of problem (1)–(2).

Theorem 2. *Let $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ be L^1 -Carathéodory functions, and let $I_{ik}, J_{ij} : [a, b] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous functions for $i = 0, 1, k = 1, 2, \dots, n$, and $j = 1, 2, \dots, m$. Then there is at least one pair of functions $(u, v) \in X$, which is a solution of (1)–(2).*

Proof. Define the operators $T_1 : X \rightarrow X_1, T_2 : X \rightarrow X_2$, and $T : X \rightarrow X$ by

$$T(u, v) = (T_1(u, v), T_2(u, v)) \tag{5}$$

with

$$\begin{aligned}
 (T_1(u, v))(x) = & A_1 + B_1(x - a) \\
 & + \sum_{x_k < x} [I_{0k}(x_k, u(x_k), u'(x_k)) + I_{1k}(x_k, u(x_k), u'(x_k))(x - x_k)] \\
 & - (x - a) \sum_{k=1}^n I_{1k}(x_k, u(x_k), u'(x_k)) \\
 & + \int_a^b G_1(x, s)f(s, u(s), u'(s), v(s), v'(s)) ds,
 \end{aligned}$$

$$\begin{aligned}
(T_2(u, v))(x) &= A_2 + \frac{B_2 - A_2}{b - a}(x - a) \\
&+ \sum_{\tau_j < x} [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j)] \\
&- \frac{x - a}{b - a} \sum_{j=1}^m [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j)] \\
&+ \int_a^b G_2(x, s)h(s, u(s), u'(s), v(s), v'(s)) \, ds,
\end{aligned}$$

where $G_1(x, s)$ and $G_2(x, s)$ are given by (3) and (4), respectively.

By Lemma 1, it is obvious that the fixed points of T are solutions of (1)–(2), so we shall prove that T has a fixed point, following, for clearness, several steps.

Step 1. T is well defined and continuous in X .

As $f, h : [a, b] \times \mathbb{R}^4 \rightarrow \mathbb{R}$ are L^1 -Carathéodory functions, then $T_1(u, v) \in PC_1^1$ and $T_2(u, v) \in PC_2^1$. In fact, $T(u, v) = (T_1(u, v), T_2(u, v))$ is continuous and

$$\begin{aligned}
(T_1(u, v))'(x) &= B_1 + \sum_{x_k < x} I_{1k}(x_k, u(x_k), u'(x_k)) - \sum_{k=1}^n I_{1k}(x_k, u(x_k), u'(x_k)) \\
&- \int_a^x f(s, u(s), u'(s), v(s), v'(s)) \, ds, \\
(T_2(u, v))'(x) &= \frac{B_2 - A_2}{b - a} + \sum_{\tau_j < x} J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) \\
&- \frac{1}{b - a} \sum_{j=1}^m [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j)] \\
&- \frac{x - a}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) \\
&+ \int_a^b \frac{\partial G_2}{\partial x}(x, s)h(s, u(s), u'(s), v(s), v'(s)) \, ds
\end{aligned}$$

with

$$\frac{\partial G_2}{\partial x}(x, s) = \frac{1}{b - a} \begin{cases} s - a, & a \leq x \leq s \leq b, \\ b - s, & a \leq s \leq x \leq b. \end{cases} \quad (6)$$

Therefore, $T_1(u, v) \in X_1$, $T_2(u, v) \in X_2$, and $T(u, v) \in X$.

Step 2. TB is uniformly bounded in $B \subseteq X$.

Let B be a bounded set of X . As f and h are L^1 -Carathéodory functions, there exists $\phi_\rho, \psi_\rho \in L^1([a, b])$ with $\rho_1 > 0$ such that

$$\max\{\|u\|_{X_1}, \|v\|_{X_2}\} < \rho_1, \tag{7}$$

we have

$$\begin{aligned} |f(x, u(x), u'(x), v(x), v'(x))| &\leq \phi_\rho(x), \\ |h(x, u(x), u'(x), v(x), v'(x))| &\leq \psi_\rho(x), \quad \text{a.e. } x \in [a, b]. \end{aligned}$$

For $(u, v) \in B$, let

$$M_1(s) := \sup_{x \in [a, b]} |G_1(x, s)|, \quad M_2(s) := \sup_{x \in [a, b]} |G_2(x, s)|. \tag{8}$$

By the continuity of functions I_{ik}, J_{ij} for $i = 0, 1, k = 1, 2, \dots, n$, and $j = 1, 2, \dots, m$, there are positive constants φ_{ik} and φ_{ij}^* such that

$$|I_{ik}(x_k, u(x_k), u'(x_k))| \leq \varphi_{ik} \quad \text{and} \quad |J_{ij}(\tau_j, v(\tau_j), v'(\tau_j))| \leq \varphi_{ij}^*.$$

Moreover,

$$\begin{aligned} &\|T_1(u, v)(x)\| \\ &\leq \sup_{x \in [a, b]} \left(|A_1| + |B_1| |x - a| \right. \\ &\quad + \sum_{x_k < x} \left[|I_{0k}(x_k, u(x_k), u'(x_k)) + I_{1k}(x_k, u(x_k), u'(x_k))(x - x_k)| \right] \\ &\quad + |(x - a)| \sum_{k=1}^n |I_{1k}(x_k, u(x_k), u'(x_k))| \\ &\quad \left. + \int_a^b |G_1(x, s)| |f(s, u(s), u'(s), v(s), v'(s))| ds \right) \\ &\leq |A_1| + |B_1|(b - a) + \sum_{k=1}^n [\varphi_{0k} + 2(b - a)\varphi_{1k}] + \int_a^b M_1(s)\phi_\rho(s) ds < +\infty, \end{aligned}$$

$$\begin{aligned} &\|(T_1(u, v))'(x)\| \\ &\leq \sup_{x \in [a, b]} \left(|B_1| + \sum_{x_k < x} |I_{1k}(x_k, u(x_k), u'(x_k))| \right. \\ &\quad + \sum_{k=1}^n |I_{1k}(x_k, u(x_k), u'(x_k))| + \int_a^b |f(s, u(s), u'(s), v(s), v'(s))| ds \left. \right) \\ &\leq |B_1| + 2 \sum_{k=1}^n \varphi_{1k} + \int_a^b \phi_\rho(s) ds < +\infty, \end{aligned}$$

$$\begin{aligned}
& \|T_2(u, v)(x)\| \\
& \leq \sup_{x \in [a, b]} \left(|A_2| + \frac{|B_2 - A_2|}{b - a} |x - a| \right. \\
& \quad + \sum_{\tau_j < x} [|J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j) |] \\
& \quad + \frac{|x - a|}{b - a} \sum_{j=1}^m |J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j) | \\
& \quad \left. + \int_a^b |G_2(x, s)| |h(s, u(s), u'(s), v(s), v'(s))| ds \right) \\
& \leq |A_2| + |B_2 - A_2| + 2 \sum_{\tau_j < t} [\varphi_{0j}^* + \varphi_{1j}^*(b - a)] + \int_a^b M_2(s) \psi_\rho(s) ds < +\infty,
\end{aligned}$$

and, by (6),

$$\begin{aligned}
& \| (T_2(u, v))'(x) \| \\
& \leq \sup_{x \in [a, b]} \left(\frac{|B_2 - A_2|}{b - a} + \sum_{\tau_j < x} |J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))| \right. \\
& \quad + \frac{1}{b - a} \sum_{j=1}^m |J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(x - \tau_j) | \\
& \quad + \frac{|x - a|}{b - a} \sum_{j=1}^m |J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))| \\
& \quad \left. + \frac{1}{b - a} \int_a^b \left| \frac{\partial G_2}{\partial x}(x, s) \right| |h(s, u(s), u'(s), v(s), v'(s))| ds \right) \\
& \leq \frac{|B_2 - A_2|}{b - a} + \frac{1}{b - a} \sum_{j=1}^m \varphi_{0j}^* + 3 \sum_{j=1}^m \varphi_{1j}^* + \frac{1}{b - a} \int_a^b \left| \frac{\partial G_2}{\partial x}(x, s) \right| \psi_\rho(s) ds < +\infty.
\end{aligned}$$

So, TB is uniformly bounded on X .

Step 3. T is equicontinuous on each interval $]x_k, x_{k+1}[\times]\tau_j, \tau_{j+1}[$, that is, T_1B is equicontinuous on each interval $]x_k, x_{k+1}[$ for $k = 0, 1, \dots, n$ with $x_0 = a$ and $x_{n+1} = b$, and T_2B is equicontinuous on each interval $]\tau_j, \tau_{j+1}[$ for $j = 0, 1, \dots, m$ with $\tau_0 = a$ and $\tau_{m+1} = b$.

Consider $J \subseteq]x_k, x_{k+1}[$ and $\iota_1, \iota_2 \in J$ such that $\iota_1 \leq \iota_2$.

So, by the continuity of G_1 ,

$$\begin{aligned} & |T_1(u, v)(\iota_1) - T_1(u, v)(\iota_2)| \\ &= \left| B_1(\iota_1 - \iota_2) + \sum_{x_k < \iota_1} [I_{0k}(x_k, u(x_k), u'(x_k)) \right. \\ &\quad \left. + I_{1k}(x_k, u(x_k), u'(x_k))(\iota_1 - x_k)] - (\iota_1 - \iota_2) \sum_{k=1}^n I_{1k}(x_k, u(x_k), u'(x_k)) \right. \\ &\quad \left. - \sum_{x_k < \iota_2} [I_{0k}(x_k, u(x_k), u'(x_k)) + I_{1k}(x_k, u(x_k), u'(x_k))(\iota_2 - x_k)] \right. \\ &\quad \left. + \int_a^b [G_1(\iota_1, s) - G_1(\iota_2, s)] f(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\ &\rightarrow 0 \quad \text{as } \iota_1 \rightarrow \iota_2, \end{aligned}$$

$$\begin{aligned} & |(T_1(u, v)(\iota_1))' - (T_1(u, v)(\iota_2))'| \\ &= \left| \sum_{x_k < \iota_1} I_{1k}(x_k, u(x_k), u'(x_k)) - \sum_{x_k < \iota_2} I_{1k}(x_k, u(x_k), u'(x_k)) \right. \\ &\quad \left. - \int_{\iota_1}^{\iota_2} f(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\ &\rightarrow 0 \quad \text{as } \iota_1 \rightarrow \iota_2, \end{aligned}$$

$$\begin{aligned} & |T_2(u, v)(\iota_1) - T_2(u, v)(\iota_2)| \\ &= \left| \frac{B_2 - A_2}{b - a}(\iota_1 - \iota_2) + \sum_{x_k < \iota_1} [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) \right. \\ &\quad \left. + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\iota_1 - \tau_j)] \right. \\ &\quad \left. + \frac{\iota_2 - \iota_1}{b - a} \sum_{j=1}^m J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) - \frac{\iota_1 - a}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\iota_1 - \tau_j) \right. \\ &\quad \left. - \sum_{x_k < \iota_2} [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\iota_2 - \tau_j)] \right. \\ &\quad \left. + \frac{\iota_2 - a}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\iota_2 - \tau_j) \right. \\ &\quad \left. + \int_a^b [G_2(\iota_1, s) - G_2(\iota_2, s)] h(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\ &\rightarrow 0 \quad \text{as } \iota_1 \rightarrow \iota_2, \end{aligned}$$

and

$$\begin{aligned}
& |(T_2(u, v)(\iota_1))' - (T_2(u, v)(\iota_2))'| \\
&= \left| \sum_{\tau_j < \iota_1} J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) - \frac{1}{b-a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\iota_1 - \iota_2) \right. \\
&\quad + \frac{(\iota_1 - \iota_2)}{b-a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) - \sum_{\tau_j < \iota_2} J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) \\
&\quad \left. + \frac{1}{b-a} \int_{\iota_1}^{\iota_2} \frac{\partial G_2}{\partial x}(x, s) h(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\
&\rightarrow 0 \quad \text{as } \iota_1 \rightarrow \iota_2,
\end{aligned}$$

and $\partial G_2/\partial x$ given by (6).

Step 4. TB is equiconvergent, that is, T_1B is equiconvergent at $x = x_k^+$ for $k = 0, 1, \dots, n$, and T_2B is equiconvergent at $\tau = \tau_j^+$ for $\tau = 1, \dots, m$.

$$\begin{aligned}
& |T_1(u, v)(x) - T_1(u, v)(x_k^+)| \\
&= \left| B_1(x - x_k^+) + \sum_{x_k < x} [I_{0k}(x_k, u(x_k), u'(x_k)) + I_{1k}(x_k, u(x_k), u'(x_k))](x - x_k) \right. \\
&\quad - (x - x_k^+) \sum_{k=1}^n I_{1k}(x_k, u(x_k), u'(x_k)) \\
&\quad - \sum_{x_k < x_k^+} [I_{0k}(x_k, u(x_k), u'(x_k)) + I_{1k}(x_k, u(x_k), u'(x_k))](x_k^+ - x_k) \\
&\quad \left. + \int_a^b [G_1(x, s) - G_1(x_k^+, s)] f(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\
&\rightarrow 0 \quad \text{uniformly as } x \rightarrow x_k^+,
\end{aligned}$$

and

$$\begin{aligned}
& |(T_1(u, v)(x))' - (T_1(u, v)(x_k^+))'| \\
&= \left| \sum_{x_k < x} I_{1k}(x_k, u(x_k), u'(x_k)) - \sum_{x_k < x_k^+} I_{1k}(x_k, u(x_k), u'(x_k)) \right. \\
&\quad \left. - \int_{x_k^+}^x f(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\
&\rightarrow 0 \quad \text{uniformly as } x \rightarrow x_k^+.
\end{aligned}$$

So, T_1B is equiconvergent at $x = x_{k+}$ for $k = 0, 1, \dots, n$.

Similarly,

$$\begin{aligned}
 & |T_2(u, v)(\tau) - T_2(u, v)(\tau_j^+)| \\
 &= \left| \frac{B_2 - A_2}{b - a} (\tau - \tau_j^+) \right. \\
 &+ \sum_{\tau_j < \tau} [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\tau - \tau_j)] \\
 &+ \frac{\tau_j^+ - \tau}{b - a} \sum_{j=1}^m J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) - \frac{\tau - a}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\tau - \tau_j) \\
 &- \sum_{\tau_j < \tau_j^+} [J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) + J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\tau_j^+ - \tau_j)] \\
 &+ \frac{\tau_j^+ - a}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\tau_j^+ - \tau_j) \\
 &+ \left. \int_a^b [G_2(\tau, s) - G_2(\tau_j^+, s)] h(s, u(s), u'(s), v(s), v'(s)) \, ds \right| \\
 &\rightarrow 0 \quad \text{uniformly as } \tau \rightarrow \tau_j^+,
 \end{aligned}$$

and

$$\begin{aligned}
 & |(T_2(u, v)(\tau))' - (T_2(u, v)(\tau_j^+))'| \\
 &= \left| \sum_{\tau_j < \tau} J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) - \frac{1}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j))(\tau - \tau_j^+) \right. \\
 &+ \frac{(\tau - \tau_j^+)}{b - a} \sum_{j=1}^m J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) - \sum_{\tau_j < \tau_j^+} J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) \\
 &+ \frac{1}{b - a} \left(\int_{\tau_j^+}^{\tau} (b - s) h(s, u(s), u'(s), v(s), v'(s)) \, ds \right. \\
 &+ \left. \int_{\tau_j^+}^{\tau} (s - a) h(s, u(s), u'(s), v(s), v'(s)) \, ds \right) \Big| \\
 &\rightarrow 0 \quad \text{uniformly as } \tau \rightarrow \tau_j^+.
 \end{aligned}$$

Then T_2B is equiconvergent at $\tau = \tau_j^+$ for $\tau = 1, \dots, m$.

Like this, T_1 and T_2 maps bounded sets into relatively compact sets, that is, $T_1 : X \rightarrow X_1$ and $T_2 : X \rightarrow X_2$ are compacts. Therefore, $T : X \rightarrow X$ is compact (for details, see [18, Lemma 2.4]).

Step 5. $T : X \rightarrow X$ has a fixed point.

In order to apply Schauder's fixed point theorem for operator $T(u, v)$, we need to prove that $TD \subset D$ for some closed, bounded, and convex $D \subset X$.

Consider

$$D := \{(u, v) \in X : \|(u, v)\|_X \leq \rho_2\}$$

with $\rho_2 > 0$ such that

$$\begin{aligned} \rho_2 := \max \left\{ \rho_1, |A_1| + |B_1|(b-a) + \sum_{k=1}^n [\varphi_{0k} + 2(b-a)\varphi_{1k}] + \int_a^b M_1(s)\phi_\rho(s) ds, \right. \\ |B_1| + 2 \sum_{k=1}^n \varphi_{1k} + \int_a^b \phi_\rho(s) ds, \\ |A_2| + |B_2 - A_2| + 2 \sum_{\tau_j < x} [\varphi_{0j}^* + \varphi_{1j}^*(b-a)] + \int_a^b M_2(s)\psi_\rho(s) ds, \\ \left. \frac{|B_2 - A_2|}{b-a} + \frac{1}{b-a} \sum_{j=1}^m \varphi_{0j}^* + 3 \sum_{j=1}^m \varphi_{1j}^* + \frac{1}{b-a} \int_a^b \left| \frac{\partial G_2}{\partial x}(x, s) \right| \psi_\rho(s) ds \right\} \end{aligned}$$

with ρ_1 given by (7) according to Step 2, and M_1, M_2 are given by (8).

Following similar arguments as in Step 2, we obtain

$$\begin{aligned} \|T(u, v)\|_X &= \|(T_1(u, v), T_2(u, v))\|_X \\ &= \max\{\|T_1(u, v)\|_{X_1}, \|T_2(u, v)\|_{X_2}\} \\ &= \max\{\|T_1(u, v)\|, \|(T_1(u, v))'\|, \|T_2(u, v)\|, \|(T_2(u, v))'\|\} \\ &\leq \rho_2 \end{aligned}$$

and $TD \subset D$.

By Schauder's fixed point theorem, the operator T given by (5) has a fixed point (u_0, v_0) . Thus, problem (1)–(2) has at least one pair solution $(u, v) \in X$. \square

4 Example

Consider the coupled system of the second-order differential equations with the mixed boundary conditions

$$\begin{aligned} u''(x) &= \operatorname{sgn}\left(x - \frac{1}{2}\right)u(x)v(x) + (u'(x))^2v'(x), \quad x \in]0, 1[, \\ v''(x) &= -(x+1)(v'(x))^2u(x) + (u'(x))^3e^{-v(x)}, \\ u(0) &= 1, \quad u'(1) = \frac{1}{2}, \\ v(0) &= 1, \quad v(1) = 2 \end{aligned} \tag{9}$$

and the generalized impulsive conditions

$$\begin{aligned} \Delta u(x_k) &= (u(x_k))^2(1 - x_k) + u'(x_k), \\ \Delta u'(x_k) &= \sum_{k=1}^3 x_k u(x_k) u'(x_k), \quad k = 1, 2, 3, \\ \Delta v(\tau_j) &= \tau_j |v(\tau_j)| v'(\tau_j), \\ \Delta v'(\tau_j) &= \sum_{j=1}^2 (1 - \tau_j) \frac{v(\tau_j)}{2} (v'(\tau_j))^2, \quad j = 1, 2, \end{aligned} \tag{10}$$

where $0 < x_1 < x_2 < x_3 < 1, 0 < \tau_1 < \tau_2 < 1$.

This problem is a particular case of system (1)–(2) with

$$\begin{aligned} f(x, \alpha, \beta, \gamma, \delta) &= \operatorname{sgn}\left(x - \frac{1}{2}\right) \alpha \beta + \gamma^2 \delta, \\ h(x, \alpha, \beta, \gamma, \delta) &= -(x + 1) \delta^2 \alpha + \gamma^3 e^{-\beta}, \\ A_1 &= 1, \quad B_1 = \frac{1}{2}, \quad A_2 = 1, \quad B_2 = 2, \\ I_{0k}(x_k, \alpha, \beta) &= \alpha^2(1 - x_k) + \gamma, \quad I_{1k}(x_k, \alpha, \beta) = \sum_{k=1}^3 x_k \alpha \beta, \\ J_{0j}(\tau_j, \gamma, \delta) &= \tau_j |\beta| \delta, \quad J_{1j}(\tau_j, \gamma, \delta) = \sum_{j=1}^2 (1 - \tau_j) \frac{\beta}{2} \delta^2. \end{aligned}$$

In fact, f, h are L^1 -Carathéodory functions with $\rho > 0$ such that

$$\max\{|\alpha|, |\beta|, |\gamma|, |\delta|\} < \rho,$$

we have

$$\begin{aligned} |f(x, \alpha, \beta, \gamma, \delta)| &\leq \rho^2 + \rho^3 := \phi_\rho(x), \\ |h(x, \alpha, \beta, \gamma, \delta)| &\leq (x + 1) \rho^3 + \rho^4 := \psi_\rho(x). \end{aligned}$$

Moreover,

$$\begin{aligned} I_{0k}(x_k, \alpha, \beta) &= \alpha^2(1 - x_k) + \gamma, \quad I_{1k}(x_k, \alpha, \beta) = \sum_{k=1}^3 x_k \alpha \beta, \\ J_{0j}(\tau_j, \gamma, \delta) &= \tau_j |\beta| \delta \quad \text{and} \quad J_{1j}(\tau_j, \gamma, \delta) = \sum_{j=1}^2 (1 - \tau_j) \frac{\beta}{2} \delta^2 \end{aligned}$$

are continuous functions in $[0, 1] \times \mathbb{R}^2$.

Therefore, by Theorem 2, problem (9)–(10) has at least one solution $(u, v) \in X$.

5 The transverse vibration system of elastically coupled double-string

Consider the transverse vibration system of elastically coupled double-string with damping. The strings have the same length L , are attached by a viscoelastic element, and stretched at a constant tension according to Fig. 1.

By [26], the system of elastically coupled double-string stationary model is given by the second-order nonlinear system of differential equations

$$\begin{aligned} S_1 u''(x) - K(u(x) - v(x)) &= -l_1(x), \\ S_2 v''(x) - K(v(x) - u(x)) &= -l_2(x), \end{aligned} \quad (11)$$

where $x \in [0, L]$,

- $u(x), v(x)$ are the transverse deflections of strings u and v , respectively;
- $l_1(x)$ and $l_2(x)$ are the exciting distributed load;
- K is the modulus of Kelvin–Voigt viscoelastic;
- S_1, S_2 are the string tensions of u and v , respectively.

Adding to system (11) the boundary conditions

$$u(0) = 0, \quad u'(L) = B_1, \quad v(0) = 0, \quad v(L) = 0, \quad (12)$$

we remark that strings u and v have different behaviors at the end points. Moreover, we consider impulsive conditions that may depend on the string deflections and on the slope of the corresponding deflections:

$$\begin{aligned} I_{0k}(x_k, u(x_k), u'(x_k)) &= \eta_1 u(x_k) + \eta_2 u'(x_k) + x_k, \\ I_{1k}(x_k, u(x_k), u'(x_k)) &= \eta_3 u(x_k) + \eta_4 u'(x_k) + x_k, \\ J_{0j}(\tau_j, v(\tau_j), v'(\tau_j)) &= \eta_5 v(\tau_j) + \eta_6 v'(\tau_j) + \tau_j, \\ J_{1j}(\tau_j, v(\tau_j), v'(\tau_j)) &= \eta_7 v(\tau_j) + \eta_8 v'(\tau_j) + \tau_j. \end{aligned} \quad (13)$$

Here $B_1, \eta_i \in \mathbb{R}$, $i = 1, \dots, 8$, $k = 1, \dots, n$, and $j = 1, \dots, m$. In this way, we have a particular case of (1)–(2).




Figure 1. Elastically coupled double-string.

In system (11)–(13), it follows that

$$f(x, \alpha, \beta, \gamma, \delta) = \frac{1}{S_1} [K(\alpha - \gamma) - l_1(x)],$$

$$h(x, \alpha, \beta, \gamma, \delta) = \frac{1}{S_2} [+K(\gamma - \alpha) - l_2(x)],$$

and

$$I_{0k}(x_k, \alpha, \gamma) = \eta_1\alpha + \eta_2\gamma, \quad I_{1k}(x_k, \alpha, \gamma) = \eta_3\alpha + \eta_4\gamma + x_k,$$

$$J_{0j}(\tau_j, \beta, \delta) = \eta_5\beta + \eta_6\delta, \quad J_{1j}(\tau_j, \beta, \delta) = \eta_7\beta + \eta_8\delta + \tau_j$$

with $k = 1, \dots, n$ and $j = 1, \dots, m$.

Notice that f, h are L^1 -Carathéodory functions in $[0, L] \times \mathbb{R}^4$ with $\rho > 0$ and

$$\max\{|\alpha|, |\beta|, |\gamma|, |\delta|\} < \rho,$$

$$|f(x, \alpha, \beta, \gamma, \delta)| \leq \frac{1}{|S_1|} [2|K|\rho + |l_1(x)|] := \phi_\rho(x),$$

$$|h(x, \alpha, \beta, \gamma, \delta)| \leq \frac{1}{|S_2|} [2|K|\rho + |l_2(x)|] := \psi_\rho(x).$$

As the impulsive conditions, I_{ik} for $i = 1, 2, k = 1, \dots, n$ and J_{ij} for $j = 1, \dots, m$ are continuous functions on $[0, L] \times \mathbb{R}^2$. Then the assumptions of Theorem 2 are satisfied, and therefore, problem (11)–(13) has at least one solution $(u, v) \in X$.

Acknowledgment. The authors thank the referees for suggestions and comments to further improve this work.

References

1. A. Abdel-Rady, A. El-Sayed, S. Rida, I. Ameen, On some impulsive differential equations, *Math. Sci. Lett.*, **1**(2):105–113, 2012.
2. M. Benchohra, J. Henderson, S. Ntouyas, *Impulsive Differential equations and Inclusions*, Hindawi Publishing Corporation, New York, 2006.
3. M. Benchohra, B. Slimani, Existence and uniqueness of solutions to impulsive fractional differential equations, *Electron. J. Differ. Equ.*, **10**:1–11, 2009.
4. A. Catllá, D. Schaeffer, T. Witeliski, E. Monson, A. Lin, On spiking models for synaptic activity and impulsive differential equations, *SIAM Rev.*, **50**(3):553–569, 2008.
5. T. Fay, S. D. Graham, Coupled spring equations, *Int. J. Math. Educ. Sci. Technol.*, **34**(1):65–79, 2003.
6. J. Fialho, F. Minhós, Fourth order impulsive periodic boundary value problems, *Differ. Equ. Dyn. Syst.*, **23**(2):117–127, 2015.
7. J. Fialho, F. Minhós, High order periodic impulsive problems, in M. de León et al. (Eds.), *Dynamical Systems, Differential Equations and Applications, AIMS Proceedings 2015*, AIMS, 2015, pp. 446–454.

8. D. Guo, Boundary value problems for impulsive integro-differential equation on unbounded domains in a Banach space, *Appl. Math. Comput.*, **99**:1–15, 1999.
9. G. Infante, F.M. Minhós, P. Pietramala, Non-negative solutions of systems of ODEs with coupled boundary conditions, *Commun. Nonlinear Sci. Numer. Simul.*, **17**(12):4952–4960, 2012.
10. G. Infante, P. Pietramala, Existence and multiplicity of non-negative solutions for systems of perturbed Hammerstein integral equations, *Nonlinear Anal., Theory Methods Appl.*, **71**(3–4): 1301–1310, 2009.
11. G. Infante, P. Pietramala, Nonnegative solutions for a system of impulsive BVPs with nonlinear nonlocal BCs, *Nonlinear Anal. Model. Control*, **19**(3):413–431, 2014.
12. T. Jankowski, First-order impulsive ordinary differential equations with advanced arguments, *J. Math. Anal. Appl.*, **331**(3):1–12, 2007.
13. X. Li, M. Bohner, C-K. Wang, Impulsive differential equations: Periodic solutions and applications, *Automatica*, **52**:173–178, 2015.
14. X. Li, S. Song, Impulsive control for existence, uniqueness and global stability of periodic solutions of recurrent neural networks with discrete and continuously distributed delays, *IEEE Trans. Neural Networks Learn. Syst.*, **24**(6):868–877, 2013.
15. X. Li, S. Song, Stabilization of delay systems: Delay dependent impulsive control, *IEEE Trans. Autom. Control*, **62**:406–411, 2017.
16. X. Li, J. Wu, Stability of nonlinear differential systems with state-dependent delayed impulses, *Automatica*, **64**:63–69, 2016.
17. Y. Liu, Existence results on positive periodic solutions for impulsive functional differential equations, *Glas. Mat., III. Ser.*, **46**(1):149–165, 2011.
18. Y. Liu, Existence of solutions of boundary value problems for coupled singular differential equations on whole line with impulses, *Mediterr. J. Math.*, **12**:697–716, 2015.
19. H. Lü, H. Yu, Y. Liu, Positive solutions for singular boundary value problems of a coupled system of differential equations, *J. Math. Anal. Appl.*, **302**(66):14–29, 2005.
20. F. Minhós, Impulsive problems on the half-line with infinite impulse moments, *Lith. Math. J.*, **57**(1):69–79, 2017.
21. F. Minhós, R. Carapinha, On higher order nonlinear impulsive boundary value problems, in M. de León et al. (Eds.), *Dynamical Systems, Differential Equations and Applications, AIMS Proceedings 2015*, AIMS, 2015, pp. 851–860.
22. F. Minhós, I. Coxe, System of coupled clamped beam equations: Existence and localization results, *Nonlinear Anal., Real World Appl.*, **35**(1):45–60, 2017.
23. F. Minhós, R. de Sousa, On the solvability of third-order three points systems of differential equations with dependence on the first derivative, *Bull. Braz. Math. Soc. (N.S.)*, **48**(3):485–503, 2017.
24. F. Minhós, R. de Sousa, Existence of solution for functional coupled systems with full nonlinear terms and applications to a coupled mass-spring model, to appear.
25. F. Minhós, J. Fialho, On the solvability of some fourth-order equations with functional boundary conditions, *Discrete Contin. Dyn. Syst.*, **2009**(Suppl.):564–573, 2009.

26. Z. Oniszczyk, Damped vibration analysis of an elastically connected complex double-string system, *J. Sound Vib.*, **264**:253–271, 2003.
27. V. Rodrigues, *Estudo das Vibrações Transversais em um Sistema Viscoelástico Acoplado de Duas Cordas*, Santa Maria, RS, Brasil, 2013.
28. K. Shah, H. Khalil, R.A. Khan, Investigation of positive solution to a coupled system of impulsive boundary value problems for nonlinear fractional order differential equations, *Chaos Solitons Fractals*, **77**:240–246, 2015.
29. I. Stamova, G. Stamov, *Applied Impulsive Mathematical Models*, Springer, 2016.
30. I. Talib, N.A. Asif, C. Tunc, Coupled lower and upper solution approach for the existence of solutions of nonlinear coupled system with nonlinear coupled boundary conditions, *Proyecciones*, **35**(1):99–117, 2016.
31. E. Zeidler, *Nonlinear Functional Analysis and Its Applications: Fixed-Point Theorems*, Springer, New York, 1986.